



University of the Pacific Scholarly Commons

Dugoni School of Dentistry Faculty Presentations

Arthur A. Dugoni School of Dentistry

3-24-2017

The Role of Neurotrophic Factors in Dental Pulp Stem Cell

Nan (Tori) Xiao

University of the Pacific, nxiao@pacific.edu

Kamille Mercado

University of the Pacific

Follow this and additional works at: <https://scholarlycommons.pacific.edu/dugoni-facpres>



Part of the [Dentistry Commons](#)

Recommended Citation

Xiao, N., & Mercado, K. (2017). The Role of Neurotrophic Factors in Dental Pulp Stem Cell. Paper presented at American Association of Dental Research San Francisco Section Research Event in San Francisco, CA.

<https://scholarlycommons.pacific.edu/dugoni-facpres/51>

This Poster is brought to you for free and open access by the Arthur A. Dugoni School of Dentistry at Scholarly Commons. It has been accepted for inclusion in Dugoni School of Dentistry Faculty Presentations by an authorized administrator of Scholarly Commons. For more information, please contact mgibney@pacific.edu.

The Role of Neurotrophic factors in Dental Pulp Stem Cell

Nan (Tori) Xiao^{1*}, Kamille Mercado²

^{1*}Department of Biomedical Sciences, ² Doctor of Dental Surgery Program (DDS 2017), University of the Pacific,
Arthur A. Dugoni School of Dentistry, San Francisco




Abstract



Neurotrophic factors are growth factors that can nourish neurons and promote neuron survival and regeneration. They have been studied as potential drug candidates for treating neurodegenerative diseases, such as Alzheimer's disease and Parkinson's disease. Since their identification, there are new evidences to indicate that neurotrophic factors are also expressed in non-neuronal tissues and promote the regeneration of tissue-specific adult stem cells.

Dental pulp stem cells (DPSCs) are multipotent mesenchymal cells that can differentiate into different cells originally derived from ectoderm, mesoderm and endoderm. Researches indicate that human DPSCs are able to differentiate into neuron, blood vessel, dentin, bone and adipose tissue under appropriate conditions. These properties make DPSCs a promising resource for stem cell therapy.

This study aims to investigate the role of neurotrophic factors in the DPSCs. The hypothesis is that neurotrophic factors promotes the DPSCs proliferation, and may also be involved in DPSCs migration and differentiation. Using Quantitative PCR, the expression of the neurotrophic family members and receptors in the DPSC were determined. Preliminary data indicated that neurotrophic factor PSPN has a dose dependent effect on the DPSCs migration. The future direction is to investigate the effect of different neurotrophic factors on DPSCs behavior, and the downstream signaling pathways involved in the process. This work will be important to understand the molecular mechanism of the dental pulp regeneration post trauma and pulpectomy, and will also shed light on the potential application of neurotrophic factors and DPSCs in other adult tissue regeneration.



Recombinant hPSPN increases DPSC migration.



Background

Neurotrophic factors include NGF, BDNF, NTs and the glial cell derived neurotrophic factors (GDNF) family. Besides GDNF, the family also includes neurturin (NRTN), artemin (ARTN) and persephin (PSPN). The GDNF family neurotrophic factors are also considered members of the transforming growth factor-beta (TGF-beta) family. By activating the GDNF family receptors (GFR) and the co-receptor tyrosine kinase RET, GDNF family is reported to improve neuron survival, proliferation and migration. GDNF is also reported enriched in salivary gland stem cells, and increases post radiation salivary gland regeneration.

Materials & Methods

Human DPSCs cells were gift from Dr. Songtao Shi, Chair and Professor Department of Anatomy & Cell Biology, University of Pennsylvania School of Dental Medicine
DPSCs mRNA was extracted from samples of five different patients, and reverse transcribed into cDNA. Primer sequences were designed with Primer3web online software. Quantitative PCR was performed on the cDNA samples using a 7900HT detection system (Applied Biosystems). All PCR reactions were carried out in triplicate. Quantification of the samples was calculated with the threshold cycle by $\Delta\Delta C_t$ method.
Immunofluorescent staining images were acquired using a Leica TCS SPE confocal microscope. Scratch assay images were acquired using a Leica DMiL inverted microscope.

Results

GDNF family Neurotrophic Factors and Receptors are expressed in DPSCs.

Future Directions

The effects of different Neurotrophic factors on DPSCs proliferation and differentiation under normal condition and inflammatory condition.
The molecular pathways downstream of the neurotrophic factors in regulating DPSCs behavior.

References

1. Liu H, Gronthos S, Shi S. Dental pulp stem cells. Methods Enzymol. 2006;419:99-113.
2. Kadar K, Kiraly M, Porcsalmy B, Molnar B, Racz GZ, Blazsek J, Kallo K, Szabo EL, Gera I, Gerber G, Varga G. Differentiation potential of stem cells from human dental origin - promise for tissue engineering. J Physiol Pharmacol. 2009;60 Suppl 7:167-75.
3. Fried K, Nosrat C, Lillesaar C, Hildebrand C. Molecular signaling and pulpal nerve development. Crit Rev Oral Biol Med. 2000;11(3):318-32.
4. Xiao N, Lin Y, Cao H, Sirjani D, Giaccia AJ, Koong AC, Kong CS, Diehn M, Le QT. Neurotrophic factor GDNF promotes survival of salivary stem cells. J Clin Invest. 2014;124(8):3364-77.
5. Liang CC, Park AY, Guan JL. In vitro scratch assay: a convenient and inexpensive method for analysis of cell migration in vitro. Nat Protoc. 2007;2(2):329-33